arXiv:gr-qc/9805042v1 12 May 1998

Symmetric hyperbolic systems for Bianchi equations

Miguel Á.G. Bonilla*
Departament de Física Fonamental, Universitat de Barcelona,
Av. Diagonal 647, E-08028 Barcelona, Spain
E-mail: mangel@ffn.ub.es

February 7, 2008

Abstract

We obtain a family of first-order symmetric hyperbolic systems for the Bianchi equations. They have only physical characteristics: the light cone and timelike hypersurfaces. In the proof of the hyperbolicity, new positivity properties of the Bel tensor are used.

PACS number: 04.20.Ex

1 Introduction

First-order symmetric hyperbolic (FOSH) systems have recently been the subject of intensive research, not only for their theoretical interest in the Cauchy problem, but also for their applications to numerical relativity [1]. Such systems constitute a powerful procedure that can be used to establish a well-posed initial-value formulation for physical systems, since most of them can be cast into this FOSH form [2] and, within this framework, theorems of existence and uniqueness of solutions are applicable [3] (see also [4]). More particularly, the Bianchi identities (investigated for a long time mainly for its usefulness in the study of the gravitational field propagation) are the basis from where some FOSH systems have been extracted (see [5] and references therein). Many of these results have been obtained by making use of the vacuum Bianchi identities, however the non-vacuum ones can also be applied to the construction of FOSH systems for the Einstein field equations [6].

In this paper a family of FOSH systems for the Bianchi equations is constructed. They may be useful in deriving new hyperbolic systems for Einstein's field equations, analogously to what is done in [6], where Bianchi identities are linked to equations for the spatial metric and connection. We begin in section 2 by considering the Bianchi equations applied to a double 2-form which does not satisfy *a priori* the additional Riemann symmetries. From these equations we extract a FOSH system of evolution equations having no unphysical characteristics. Actually, their only characteristics are the null cone and timelike hypersurfaces. In section 3 we briefly investigate the set of constraints that completes the hyperbolic system found in section 2. Both together, the evolution equations and the constraints, are equivalent to the original Bianchi equations. Finally, in section 4 we show that the full Riemann symmetries (together with the Einstein field equations) are conserved under the evolution. In the appendix we prove new positivity properties of the Bel tensor that are needed for the proof of the hyperbolicity.

---

*Also at Laboratori de Física Matemàtica, Societat Catalana de Física, IEC, Barcelona

2 Hyperbolizations of Bianchi equations

We recall that the Bianchi identities for the Riemann tensor read:

$$\nabla_{[\nu} R_{\alpha\beta]\lambda\mu} = 0\,, \tag{1}$$

from where it follows (by contracting, using the Riemann symmetries and the Einstein field equations with $8\pi G = c = 1$):

$$\nabla^\beta R_{\alpha\beta\lambda\mu} = 2\nabla_{[\mu}\left(T_{\lambda]\alpha} - \frac{1}{2} T g_{\lambda]\alpha}\right) \equiv J_{\alpha\lambda\mu}\,, \tag{2}$$

$T_{\alpha\beta}$ being the energy-momentum tensor. For the system (1,2), called the superior order field equations by Lichnerowicz, Einstein's field equations can be considered as initial data [7]: if they hold at the initial spatial hypersurface S, they also hold in a neighborhood of S. In the last section we will see that the symmetries can also be taken as initial data.

Consider now the set of equations (1,2), which will be called *Bianchi equations* in what follows, applied to a double 2-form $K_{\alpha\beta\lambda\mu} = K_{[\alpha\beta][\lambda\mu]}$. We do not suppose it to have the additional Riemann symmetries $K_{[\alpha\beta\lambda]\mu} = 0$ at the present moment. The Bianchi equations for $K_{\alpha\beta\lambda\mu}$ are then:

$$\nabla^\beta K_{\alpha\beta\lambda\mu} = J_{\alpha\lambda\mu}\,, \tag{3}$$

$$\nabla_{[\nu} K_{\alpha\beta]\lambda\mu} = 0\,. \tag{4}$$

This is an overdetermined linear first-order system of 48 equations for 36 unknowns which is not manifestly symmetric hyperbolic. To handle more easily equations (3,4), it is better to write them in the following compact form (in a similar way to what is done in [2] for other systems):

$$A_I^{\epsilon\ \gamma\delta\sigma\rho} \nabla_\epsilon K_{\gamma\delta\sigma\rho} = j_I\,, \tag{5}$$

where the index I lives in the space of equations, $I = (\alpha[\lambda\mu], [\nu\alpha\beta][\lambda\mu])$, and

$$A_I^{\epsilon\ \gamma\delta\sigma\rho} = \left(\delta^{\epsilon[\delta}\delta_\alpha^{\gamma]}\delta_{[\lambda}^{[\sigma}\delta_{\mu]}^{\rho]}, \delta^\epsilon_{[\nu}\delta_\alpha^{[\gamma}\delta_{\beta]}^{\delta]}\delta_{[\lambda}^{[\sigma}\delta_{\mu]}^{\rho]}\right)\,, \quad j_I = (J_{\alpha\lambda\mu}, 0)\,.$$

A tensor $H^{I\ \gamma'\delta'\sigma'\rho'}$ is said to be a *hyperbolization* of (5) if the matrix $Q^{\epsilon\ \{\gamma'\delta'\sigma'\rho'\}\{\gamma\delta\sigma\rho\}} \equiv H^{I\ \gamma'\delta'\sigma'\rho'} A_I^{\epsilon\ \gamma\delta\sigma\rho}$ acting over the space of double 2-forms satisfies the following properties: *i)* it is symmetric, and *ii)* there exists a 1-form n_ϵ such that $n_\epsilon \mathbf{Q}^\epsilon$ is positive-definite. By means of such a hyperbolization, we can transform (5) into:

$$H^{I\ \gamma\delta\sigma\rho} A_I^{\epsilon\ \gamma'\delta'\sigma'\rho'} \nabla_\epsilon K_{\gamma'\delta'\sigma'\rho'} = H^{I\ \gamma\delta\sigma\rho} j_I \,,$$

which is now a FOSH system and, therefore, it admits the theorems of existence and uniqueness of solutions.

The set of 1-forms n_ϵ satisfying property *ii)* has a clear physical meaning (see [2] for instance) related to the propagation of the physical fields involved and to the causal character of the system. In fact, a FOSH system is said to be *causal* if every n_ϵ satisfying *ii)* is a future-directed 1-form.

For the Bianchi equations (5) the following family of tensors are hyperbolizations:

$$\begin{aligned} H^{I\ \gamma'\delta'\sigma'\rho'} &= \left(\delta^{\alpha[\gamma'} t^{\delta']} h^{[\lambda\mu][\sigma'\rho']}, -\frac{3}{2} t^{[\nu} \delta^{\alpha\,|[\gamma'} \delta^{\delta']|\,\beta]} h^{[\lambda\mu][\sigma'\rho']} \right) \,, \\ h^{\lambda\mu\sigma'\rho'} &= g^{\lambda\sigma'} \left[u^{(\mu} v^{\rho')} - \frac{1}{4} \left(u_\nu v^\nu \right) g^{\mu\rho'} \right] \,, \end{aligned}$$

where $\vec{t}$, $\vec{u}$ and $\vec{v}$ are arbitrary timelike future-directed vectors. To check property *i)* we compute, for any two double 2-forms $S_{\gamma\delta\sigma\rho}$ and $T_{\gamma\delta\sigma\rho}$, the quantity

$$\begin{aligned} & Q^{\epsilon\ \{\gamma'\delta'\sigma'\rho'\}\{\gamma\delta\sigma\rho\}} S_{\gamma'\delta'\sigma'\rho'} T_{\gamma\delta\sigma\rho} = \\ &= \left[\frac{1}{2} \left(S^{\alpha\ \beta}_{\ \nu\ \sigma} T_{\alpha\mu\beta\rho} + T^{\alpha\ \beta}_{\ \nu\ \sigma} S_{\alpha\mu\beta\rho} + S^{\alpha\ \beta}_{\ \nu\ \rho} T_{\alpha\mu\beta\sigma} + T^{\alpha\ \beta}_{\ \nu\ \rho} S_{\alpha\mu\beta\sigma} \right) - \right. \\ & -\frac{1}{4} g_{\nu\mu} \left(S^{\alpha\beta\gamma}{}_\sigma T_{\alpha\beta\gamma\rho} + T^{\alpha\beta\gamma}{}_\sigma S_{\alpha\beta\gamma\rho} \right) - \frac{1}{4} g_{\sigma\rho} \left(S_\nu{}^{\alpha\beta\gamma} T_{\mu\alpha\beta\gamma} + T_\nu{}^{\alpha\beta\gamma} S_{\mu\alpha\beta\gamma} \right) + \\ & \left. + \frac{1}{8} g_{\nu\mu} g_{\sigma\rho} S_{\alpha\beta\gamma\delta} T^{\alpha\beta\gamma\delta} \right] g^{\nu\epsilon} t^\mu u^\sigma v^\rho = Q^{\epsilon\ \{\gamma'\delta'\sigma'\rho'\}\{\gamma\delta\sigma\rho\}} T_{\gamma'\delta'\sigma'\rho'} S_{\gamma\delta\sigma\rho} \,, \end{aligned}$$

from where the symmetry is clear. Property *ii)* is demonstrated in the appendix by proving that $n_\epsilon \mathbf{Q}^\epsilon$ is a positive-definite quadratic form for any n_ϵ timelike future-directed 1-form. This also shows that the system is causal.

Let us now study the characteristics of the hyperbolized Bianchi equations found above, which finally read:

$$h_{\alpha\beta}{}^{[\sigma\rho]} \left(t^{[\delta} \nabla_\nu K^{\gamma]\nu\alpha\beta} - \frac{3}{2} t_\epsilon \nabla^{[\epsilon} K^{\gamma\delta]\alpha\beta} \right) = h_{\alpha\beta}{}^{[\sigma\rho]} \left(t^{[\delta} J^{\gamma]\alpha\beta} \right) \,. \tag{6}$$

Recall that a 1-form ζ_α is, by definition, *characteristic* for the system (6) if $\det(\zeta_\epsilon \mathbf{Q}^\epsilon) = 0$. To study the kernel of the characteristic matrix, first notice that $h_{[\alpha\beta][\lambda\mu]}$ (considered as a matrix acting over 2-forms) has non-zero determinant, that is:

$$h_{[\alpha\beta][\lambda\mu]} A^{\lambda\mu} = 0 \implies A^{[\lambda\mu]} = 0 \,. \tag{7}$$

This can be easily proven by contracting (7) successively with $\vec{u}$ and $\vec{v}$. Therefore,

$$Q^{\epsilon\ \{\gamma'\delta'\sigma'\rho'\}\{\gamma\delta\sigma\rho\}}\zeta_\epsilon K_{\gamma\delta\sigma\rho} = 0 \iff t^{[\delta}\zeta_\nu K^{\gamma]\nu\alpha\beta} - \frac{3}{2} t_\epsilon \zeta^{[\epsilon} K^{\gamma\delta]\alpha\beta} = 0\,.$$

The non-trivial solutions of this system are found to be of two types: *i)* $\zeta_\alpha\zeta^\alpha = 0$ and *ii)* $\zeta_\alpha t^\alpha = 0$. In the first case, the solutions are the double 2-forms $K_{\alpha\beta\lambda\mu}$ satisfying $\zeta^\alpha K_{\alpha\beta\lambda\mu} = 0$ and $\zeta_{[\nu}K_{\alpha\beta]\lambda\mu} = 0$. In the second one the solutions are those $K_{\alpha\beta\lambda\mu}$ having in its two first indices the structure $A\,(t\wedge\zeta) + B*(t\wedge\zeta)$ ("$*$" is the usual dual operator). Therefore, as remarked above, (6) has only physical characteristics: *i)* the light cone and *ii)* timelike hypersurfaces.

3 Equivalence with the original system

From the Bianchi equations we have constructed the 36 propagation equations (6) (in the sense that they contain derivatives in any timelike direction n_α). In this section we will show that these equations, when completed with the constraints, are equivalent to the original system (3,4). Taking n_ν as the normal to an initial-value spatial hypersurface S and $\eta^{\epsilon\alpha\beta\nu}$ the spacetime volume 4-form, the constraints are:

$$n^\alpha\left(\nabla^\beta K_{\alpha\beta\lambda\mu} - J_{\alpha\lambda\mu}\right) = 0\,, \tag{8}$$

$$n_\nu \eta^{\epsilon\alpha\beta\nu}\nabla_\epsilon K_{\alpha\beta\lambda\mu} = 0\,, \tag{9}$$

where all the derivatives appearing in these expressions are tangent to S. This set of constraints is complete (there are 12 equations) and also integrable provided that $\nabla^\alpha J_{\alpha\lambda\mu} = 0$, which holds because of the Einstein field equations.

Taking into account (7), equation (6) immediately implies

$$t^{[\delta}\nabla_\nu K^{\gamma]\nu\alpha\beta} - \frac{3}{2} t_\epsilon \nabla^{[\epsilon} K^{\gamma\delta]\alpha\beta} - t^{[\delta}J^{\gamma]\alpha\beta} = 0\,. \tag{10}$$

If we multiply the previous expression by t_δ, we obtain $P^\sigma_\alpha\left(\nabla^\beta K_{\sigma\beta\lambda\mu} - J_{\sigma\lambda\mu}\right) = 0$, where P^σ_α is the orthogonal projector to t_α. Taking into account (8) and the fact that $t_\alpha n^\alpha \neq 0$, we obtain equation (3). Equation (4) is achieved in a similar way: we first multiply (10) by $\eta_{\gamma\delta\sigma\rho}t^\rho$ obtaining $P^\sigma_\nu \eta^{\epsilon\alpha\beta\nu}\nabla_\epsilon K_{\alpha\beta\lambda\mu} = 0$, and then we follow the same reasoning as before, but now making use of constraint (9). This shows that the set of equations (6,8,9) is equivalent to (3,4).

4 Riemann symmetries are conserved

Until now, we have only dealt with double 2-forms, but, actually, we are interested in the Riemann tensor. Our purpose now is to show that Bianchi equations preserve the full Riemann symmetries. The complete set of Riemann symmetries is $K_{\alpha\beta\lambda\mu} =$

$K_{[\alpha\beta][\lambda\mu]}$ and $K_{[\alpha\beta\lambda]\mu}=0$. In order to construct an appropriate system of differential equations, let us set the following definitions:

$$\Xi_{\alpha\beta\lambda\mu}\equiv K_{[\alpha\beta\lambda]\mu}\,,\ \ \Sigma_{\alpha\beta}\equiv K_{\alpha\beta}-\left(T_{\alpha\beta}-\frac{1}{2}Tg_{\alpha\beta}\right)\,,$$

where $K_{\alpha\beta}\equiv K^{\sigma}{}_{\alpha\sigma\beta}$. Then we extract, from equations (3,4) and the energy-momentum conservation equation $\nabla^{\beta}T_{\alpha\beta}=0$, the following system for the previous defined variables $\Xi_{\alpha\beta\lambda\mu}$ and $\Sigma_{\alpha\beta}$:

$$\begin{aligned}\nabla^{\alpha}\Xi_{\alpha\beta\lambda\mu}+\frac{2}{3}\nabla_{[\beta}\Sigma_{\lambda]\mu}&=0\,,\\ \nabla^{\alpha}\Sigma_{\alpha\beta}&=0\,,\\ \nabla^{\beta}\Sigma_{\alpha\beta}-\frac{1}{2}\nabla_{\alpha}\Sigma&=0\,.\end{aligned}\tag{11}$$

These are 32 equations for 32 unknowns forming a linear first-order homogeneous system whose only characteristic is the null cone. Now, we suppose that initially the double 2-form $K_{\alpha\beta\lambda\mu}$ satisfies the full Riemann symmetries ($\Xi_{\alpha\beta\lambda\mu}=0$) and that, in addition, $K_{\alpha\beta}$ satisfies Einstein's equations in the sense $\Sigma_{\alpha\beta}=0$. These initial conditions have no other solution but $\Xi_{\alpha\beta\lambda\mu}=\Sigma_{\alpha\beta}=0$ (see [4], for instance) and, therefore, the Riemann symmetries, likewise the Einstein field equations, are preserved by the Bianchi equations.

Acknowledgments

We wish to thank Carlos F. Sopuerta and Sergi R. Hildebrandt for a careful reading of this manuscript. We also acknowledge the *Comissionat per a Universitats i Recerca de la Generalitat de Catalunya* for financial support.

A Positivity property of Bel tensor

To prove that $n_{\epsilon}\mathbf{Q}^{\epsilon}$ is a positive-definite quadratic form we will make use of the standard spinor calculus [8]. First, taking into account the symmetries of $K_{\alpha\beta\lambda\mu}$, we decompose it as follows:

$$\begin{aligned}K_{\nu\mu\sigma\rho}=&\ \chi_{NMSR}\epsilon_{N'M'}\epsilon_{S'R'}+\Psi_{NMS'R'}\epsilon_{N'M'}\epsilon_{SR}\\ &+\overline{\chi}_{N'M'S'R'}\epsilon_{NM}\epsilon_{SR}+\overline{\Psi}_{N'M'SR}\epsilon_{NM}\epsilon_{S'R'}\,,\end{aligned}$$

where

$$\chi_{NMSR}=\chi_{(NM)(SR)}\,,\Psi_{NMS'R'}=\Psi_{(NM)(S'R')}\,.$$

Now we proceed to calculate

$$n_\epsilon Q^{\epsilon\ \{\gamma'\delta'\sigma'\rho'\}\{\gamma\delta\sigma\rho\}} K_{\gamma\delta\sigma\rho} K_{\gamma'\delta'\sigma'\rho'} = \Big(K^{\alpha\ \beta}_{\ \nu\ \ \sigma} K_{\alpha\mu\beta\rho} + K^{\alpha\ \beta}_{\ \nu\ \ \rho} K_{\alpha\mu\beta\sigma} - \\ -\frac{1}{2} g_{\nu\mu} K^{\alpha\beta\gamma}{}_{\sigma} K_{\alpha\beta\gamma\rho} - \frac{1}{2} g_{\sigma\rho} K_{\nu}{}^{\alpha\beta\gamma} K_{\mu\alpha\beta\gamma} + \frac{1}{8} g_{\nu\mu} g_{\sigma\rho} K_{\alpha\beta\gamma\delta} K^{\alpha\beta\gamma\delta} \Big) n^\nu t^\mu u^\sigma v^\rho \,, \tag{12}$$

which is the Bel tensor [9] for the double 2-form $K_{\alpha\beta\gamma\delta}$ contracted with four timelike future-directed vectors, $\vec{n}$, $\vec{t}$, $\vec{u}$ and $\vec{v}$. In fact, this is a little more general since we have not required symmetry between indices $\alpha\beta$ and $\lambda\mu$. In order to express (12) in spinor form it is more convenient to rewrite it in the following form:

$$\frac{1}{2} \Big(K^{\alpha\ \beta}_{\ \nu\ \ \sigma} K_{\alpha\mu\beta\rho} + K{*}^{\alpha\ \beta}_{\ \nu\ \ \sigma} K{*}_{\alpha\mu\beta\rho} + \\ + {*}K^{\alpha\ \beta}_{\ \nu\ \ \sigma} {*}K_{\alpha\mu\beta\rho} + {*}K{*}^{\alpha\ \beta}_{\ \nu\ \ \sigma} {*}K{*}_{\alpha\mu\beta\rho} \Big) n^\nu t^\mu u^\sigma v^\rho \,,$$

and now, the dual terms appearing in this expression have simple spinor expressions (see [8]). Performing the calculations we finally obtain

$$n_\epsilon Q^{\epsilon\ \{\gamma\delta\sigma\rho\}\{\gamma'\delta'\sigma'\rho'\}} K_{\gamma\delta\sigma\rho} K_{\gamma'\delta'\sigma'\rho'} = \\ = 4 \Big(\chi_{NMSR} \overline{\chi}_{N'M'S'R'} + \Psi_{NMS'R'} \overline{\Psi}_{N'M'SR} \Big) n^\nu t^\mu u^\sigma v^\rho \,, \tag{13}$$

which is always positive, and zero if and only if $\chi_{NMSR} = \Psi_{NMS'R'} = 0$. This is proven by using usual methods (see, for instance, [8, 10]): let us consider a spinor basis (ι_A, o_A) and its two associated null future-directed vectors, $l_\alpha = \iota_A \overline{\iota}_{A'}$ and $k_\alpha = o_A \overline{o}_{A'}$. Given a timelike future-pointing vector $\vec{z}$ and a future-directed null vector $\vec{x}$, there exists another future-directed null vector $\vec{y}$ such that:

$$\vec{z} = a\vec{x} + \vec{y} = a x_A \overline{x}_{A'} + y_A \overline{y}_{A'} \,, \ \ a > 0.$$

Applying this decomposition to $\vec{n}$, $\vec{t}$, $\vec{u}$ and $\vec{v}$, (13) becomes a sum of positive terms, so the positivity is shown. Suppose now that the expression (13) is zero. We can decompose the four previous vectors in many differents ways by choosing their corresponding vector $\vec{x}$ to be $\vec{l}$ or $\vec{k}$. Then, for every choice we do, among the positive terms of the sum we will get a component of χ_{NMSR} and $\Psi_{NMS'R'}$ in the spinor basis mentioned above. Therefore, making all the possible choices, we will obtain all the components zero, which means that $K_{\alpha\beta\lambda\mu} = 0$.

References

[1] Bona C, Masso J, Seidel E and Stela J 1995 A new formalism for numerical relativity *Phys. Rev. Lett.* **75** 600-3

[2] Geroch R 1995 Partial Differential Equations of Physics, in *General Relativity* ed G S Hall and J R Pulham (SUSSP Publications and IoP) *Preprint* gr-qc/9602055

[3] Friedrichs K O 1954 Symmetric hyperbolic linear differential equations *Commun. Pure Appl. Math.* **7** 345-92

[4] John F 1982 *Partial Differential Equations* (New York: Springer)

[5] Friedrich H 1996 Hyperbolic reductions for Einstein's equations *Class. Quantum Grav.* **13** 1451-69

[6] Anderson A, Choquet-Bruhat Y and York J W Jr 1997 Einstein-Bianchi Hyperbolic System for General Relativity (to appear in *Topological Methods in Nonlinear Analysis*) *Preprint* gr-qc/9710041

[7] Lichnerowicz A 1960 Ondes et radiations électromagnétiques et gravitationelles en relativité générale *Annali di matematica pura ed applicata* **50** 1-96

[8] Penrose R and Rindler W 1986 *Spinors and Space-Time* (Cambridge: Cambridge University Press)

[9] Bel L 1959 Introduction d'un tenseur du quatrième ordre *C.R. Acad. Sci.* **248** 1297-1300

[10] Bergqvist G 1998 Positivity properties of the Bel-Robinson tensor *J. Math. Phys.* **39** 2141-7